

Zeroth Semester Examination, Ides March, 709 AUC

MATH2100

ADVANCED MATHEMATICAL ANALYSIS

(Unit Courses, Inf. Tech.)

Time: TWO Hours for working

Ten minutes for perusal before examination begins

Check that this examination paper has 28 printed pages!**CREDIT WILL BE GIVEN ONLY FOR WORK WRITTEN ON
THIS EXAMINATION PAPER!**Answer as many questions as time permits, and make sure to answer AT LEAST
ONE of Questions A1, A2, A3 and AT LEAST ONE of Questions B1, B2, B3.

It is expected that AT LEAST THREE questions IN TOTAL will be attempted.

All questions carry the same number of marks.

Pocket calculators without ASCII capabilities may be used.

TRIAL EXAMINATION

FAMILY NAME (PRINT): _____

GIVEN NAMES (PRINT): _____

STUDENT NUMBER:

--	--	--	--	--	--	--	--

SIGNATURE: _____

EXAMINER'S USE ONLY			
QUESTION	MARK	QUESTION	MARK
A1		B1	
A2		B2	
A3		B3	
TOTAL MARKS			

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

PART A

Q A1. (a) Find the general solution of the system

$$\underset{\sim}{y}'(t) = A \underset{\sim}{y}(t), \quad A = \begin{bmatrix} -1 & -1 \\ 5 & 1 \end{bmatrix}.$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q A1. (a) **Working space only**

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q A1. (b) Write the general solution in real form.

(c) Identify the type and stability of the equilibrium (critical) point at the origin.

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

- Q A1. (d) Determine the slopes of trajectories where they cross the lines $y_2 = y_1$, $y_2 = -y_1$, $y_2 = -5y_1$ and $y_1 = 0$. Determine the directions of trajectories where they cross the line $y_1 = 0$. Use your results to help sketch some trajectories.

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q A2. (a) Rewrite

$$y''(t) - 2y(t) + y(t)^2 + y(t)^3 = 0$$

as a system of two coupled ordinary differential equations, and locate the equilibrium points of the system.

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q A2. (b) Determine the type and stability of the equilibrium points by linearization.

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

- Q A2. (c) Use the Second Shifting Theorem (see Table) to find the Inverse Laplace Transform of

$$F(s) = \frac{e^{-3s}}{s^3}.$$

Sketch the resulting inverse function, $f(t)$.

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q A3. (a) Using the Table provided,

- (i) Representing the hyperbolic functions in terms of exponential functions, and applying the First Shifting Theorem (see Table), show that

$$\mathcal{L}(\cosh t \sin t) = \frac{s^2 + 2}{s^4 + 4}.$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

- Q A3. (a) (ii) Use the Convolution Theorem (see Table) with $F(s) = G(s) = 1/(s + 4)$ to show that the Inverse Laplace Transform of $1/(s + 4)^2$ is

$$f(t) = te^{-4t}$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q A3. (b) Use the method of Laplace Transforms to solve the equation

$$y''(t) + 3y'(t) - 4y(t) = -5e^{-4t}$$

with Initial Conditions $y(0) = 1$, $y'(0) = -8$.

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q A3. (b) **Working space only.**

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

PART B

- Q B1. (a) For times $t > 0$, a cylinder of iron of length L , with its curved sides insulated, lies along the X -axis with its planar face at $x = 0$ maintained at constant temperature u_0 and its planar face at $x = L$ maintained at constant temperature u_1 . At time $t = 0$, the temperature distribution in the cylinder is a function $f(x)$, for $0 < x < L$. Write down the PDE, BCs and IC satisfied by the temperature $u(x, t)$ for $0 < x < L$ and $t > 0$, and deduce that the steady-state temperature $u^{SS}(x)$ which will be approached as $t \rightarrow \infty$ is

$$u^{SS}(x) = \left(\frac{u_1 - u_0}{L} \right) x + u_0 .$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

- Q B1. (b) Write $\hat{u}(x, t) = u(x, t) - u^{SS}(x)$, and find the PDE, BCs and IC satisfied by $\hat{u}(x, t)$. Work carefully through Fourier's Method of Separation of Variables and Superposition to obtain the solution $\hat{u}(x, t)$ of these equations in the case $f(x) = u_2$ (const.), and hence deduce that in this case

$$u(x, t) = \left(\frac{u_1 - u_0}{L} \right) x + u_0 + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{[u_2 - u_0 + (u_1 - u_2)(-1)^n]}{n} \sin\left(\frac{n\pi x}{L}\right) e^{-(n\pi c/L)^2 t}.$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B1. (b) **Working space only.**

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B1. (b) **Working space only.**

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B1. (b) **Working space only.**

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B2. (a) Use the formulas

$$\sin(A) \sin(B) = \frac{1}{2} [\cos(A-B) - \cos(A+B)]; \quad \sin(A) \cos(B) = \frac{1}{2} [\sin(A+B) + \sin(A-B)]$$

to deduce that if $n \neq 1$, then

$$\begin{aligned} \int \sin(nx) \sin(x) dx &= \frac{1}{2} \left(\frac{1}{n-1} \sin[(n-1)x] - \frac{1}{n+1} \sin[(n+1)x] \right) \\ \int \cos(nx) \sin(x) dx &= \frac{1}{2} \left(\frac{1}{n-1} \cos[(n-1)x] - \frac{1}{n+1} \cos[(n+1)x] \right), \end{aligned}$$

and that

$$\int \sin^2(x) dx = \frac{1}{2} \left(x - \frac{1}{2} \sin[2x] \right); \quad \int \cos(x) \sin(x) dx = -\frac{1}{4} \cos(2x).$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B2.

(b) Show that the Fourier Series corresponding to the function defined by

$$\begin{aligned} f(x) = 0, & \quad -\pi < x < 0; & f(x) = \sin(x), & \quad 0 < x < \pi; \\ \text{and} & & f(x + 2\pi) = f(x), & \quad -\infty < x < \infty, \end{aligned}$$

is

$$\frac{1}{\pi} + \frac{1}{2} \sin(x) - \frac{2}{\pi} \left(\frac{1}{(1)(3)} \cos(2x) + \frac{1}{(3)(5)} \cos(4x) + \cdots \right).$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B2. (b) **Working space only**

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B2. (b) **Working space only**

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

- Q B2. (c) By considering the value to which the series should converge at $x = \pi/4$, deduce that

$$\frac{1}{(3)(5)} - \frac{1}{(7)(9)} + \frac{1}{(11)(13)} \cdots = \frac{\pi}{4\sqrt{2}} - \frac{1}{2}.$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B3. (a) Show that the function $G(x - y, t)$ defined by

$$G(x - y, t) = \frac{1}{\sqrt{4\pi c^2 t}} e^{-(x-y)^2/(4c^2 t)}$$

satisfies the 1-dimensional Heat Equation:

$$G_t(x - y, t) = c^2 G_{xx}(x - y, t), \quad -\infty < x < \infty, \quad t > 0.$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B3. (a) **Working space only**

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B3. (b) You are given (**no need to check**) that $u(x, t)$ defined by

$$u(x, t) = \int_{-\infty}^{\infty} G(x - y, t) f(y) dy$$

satisfies the 1-dimensional Heat Equation for $-\infty < x < \infty$ and $t > 0$, and also the initial condition

$$\lim_{t \rightarrow 0^+} u(x, t) = f(x), \quad -\infty < x < \infty.$$

Show that in the case when $f(x) = u_0$ (const.) for $a < x < b$ and $f(x) = 0$ otherwise, this gives

$$u(x, t) = \frac{1}{2} u_0 \left(\operatorname{erf} \left(\frac{x - a}{\sqrt{4c^2 t}} \right) - \operatorname{erf} \left(\frac{x - b}{\sqrt{4c^2 t}} \right) \right),$$

where

$$\operatorname{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-v^2} dv.$$

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B3. (b) **Working space only**

MATH2100 — ADVANCED MATHEMATICAL ANALYSIS
Zeroth Semester Examination, Ides March, 709 AUC (continued)

Q B3.

- (c) Check that there is no flux of heat in the x -direction at the point $x = (a+b)/2$.