


P1


a)



b)



c)



(P2) Put $z = x + iy$.

2/5

a) $\bar{z} = z \Leftrightarrow x + iy = x - iy \Leftrightarrow 2iy = 0 \Leftrightarrow y = 0$
So, $\{z : \operatorname{Im}(z) = 0\}$.

b) $\bar{z} + z = 0 \Leftrightarrow 2x = 0 \Leftrightarrow x = 0$
So, $\{z : \operatorname{Re}(z) = 0\}$.

c) $\bar{z} = 4/z \Leftrightarrow z\bar{z} = 4 \Leftrightarrow |z|^2 = 4 \Leftrightarrow |z| = 2$.
So, $\{z : |z| = 2\}$.

(P3) a)
$$\frac{i}{(1-i)} + \frac{1-i}{i} = \frac{i^2 + (1-i)^2}{i(1-i)} = \frac{-1 - 2i}{1+i} \cdot \frac{1-i}{1-i}$$
$$= \frac{-1 + i - 2i - 2}{2} = \frac{-(3+i)}{2} = -\frac{3}{2} - i\frac{1}{2}$$

b) $8i = 2^3 \exp(i\pi/2)$, so cube roots are
 $\{2 \exp[i(\frac{\pi}{6} + \frac{2k\pi}{3})], k = 0, 1, 2\}$
 $= \{2 \exp^{i\pi/6}, 2 \exp^{5\pi i/6}, 2 \exp^{9\pi i/6}\}$
 $= \{2(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6}), 2(\cos \frac{5\pi}{6} + i \sin \frac{5\pi}{6}),$
 $2(\cos \frac{9\pi}{6} + i \sin \frac{9\pi}{6})\}$
 $= \{\sqrt{3} + i, -\sqrt{3} + i, -2i\}$

c) $\left(\frac{i+1}{\sqrt{2}}\right)^{1337} = \left(\exp \frac{i\pi}{4}\right)^{1337} = \exp \frac{1337\pi i}{4}$
 $= \exp \frac{\pi i}{4} = \frac{i+1}{\sqrt{2}}$

(P4) a) Given $p, q, r, s \in \mathbb{Q}$, wts:

① $(p+q\sqrt{5}) + (r+s\sqrt{5}) \in \mathbb{Q}(\sqrt{5})$ (closure under +);

② $(p+q\sqrt{5})(r+s\sqrt{5}) \in \mathbb{Q}(\sqrt{5})$ (closure under $\cdot$).

①: LHS = $(p+r) + (q+s)\sqrt{5}$, since $\mathbb{R}$ is a field,

$\Delta p+r \in \mathbb{Q}, q+s \in \mathbb{Q}$ since $\mathbb{Q}$ is a field.

This shows ①.

②: LHS = $(pr+5sq) + (r_2+ps)\sqrt{5}$, since $\mathbb{R}$

is a field, & $pr+5sq \in \mathbb{Q}, r_2+ps \in \mathbb{Q}$ since

$\mathbb{Q}$ is a field. This shows ②.

NEXT SHOW (F1)-(F7)

(F1) follows since $\mathbb{R}$ is a field.

(F2) (i) follows since $\mathbb{R}$ is a field, with $0 = 0 + 0\sqrt{5} \in \mathbb{Q}(\sqrt{5})$.

(F2) (ii) same: $-(p+q\sqrt{5}) = (-p) + (-q)\sqrt{5} \in \mathbb{Q}(\sqrt{5})$

(F4) follows since $\mathbb{R}$ is a field.

(F5) (i) same: $1 = 1 + 0\sqrt{5} \in \mathbb{Q}(\sqrt{5})$.

(ii) For $p, q \in \mathbb{Q}$ not both 0, we have

$$\frac{1}{p+q\sqrt{5}} = \frac{1}{p+q\sqrt{5}} \cdot \frac{p-q\sqrt{5}}{p-q\sqrt{5}} = \frac{p}{p^2-5q^2} + \left(\frac{-q}{p^2-5q^2}\right)\sqrt{5}$$

which is in $\mathbb{Q}(\sqrt{5})$; note in particular

$p^2 - 5q^2 \neq 0$ for $p, q \in \mathbb{Q}$, since $\sqrt{5} \notin \mathbb{Q}$.
 (F6) & (F7) follow since $\mathbb{R}$ is a field.
 Hence $\mathbb{Q}(\sqrt{5})$ is a field.

b) Suppose $\sqrt{2} = a + b\sqrt{5}$, $a, b \in \mathbb{Q}$.

Note $b \neq 0$. (If $b = 0$, then $\sqrt{2} = a \in \mathbb{Q}$, but $\sqrt{2} \notin \mathbb{Q}$).

Hence, we can rewrite $(*)$ as

$$\sqrt{2} - b\sqrt{5} = a, \text{ \& on squaring,}$$

$$5b^2 + 2 - 2\sqrt{2}\sqrt{5}b = a^2$$

$$\text{i.e., } 2\sqrt{10}b = 2 - a^2 + 5b^2$$

$$\text{since } b \neq 0, \text{ this yields } \sqrt{10} = \frac{2 - a^2 + 5b^2}{2b}, \text{ i.e.,}$$

$$\sqrt{10} \in \mathbb{Q}, \quad \text{C!}$$

assume $\mathbb{C}$ is well ordered.

(p5) As per hint: (note $-1 \notin P$, because if it were, then $(-1)(-1) \in P$ (P closed under multiplication), i.e., $1 \in P$, so both $1 \in P$ & $-1 \in P$, which is impossible).

So if $i \in P$, then $(i)(i) = -1 \in P$ (again, by closure of P under $\times$), which it isn't.

But if $-i \in P$, then $(-i)(-i) = -1 \in P$, which it isn't.

Since $i \neq 0$, we thus have a contradiction.

□