



THE UNIVERSITY
OF QUEENSLAND
AUSTRALIA

Analysis of Engineering and Scientific Data

Semester 1 – 2019

Sabrina Streipert

s.streipert@uq.edu.au

Changes in this week's tutorials

1. P02: Monday: 3-4pm: 47A-249
2. P06: Wednesday: 3-4pm: 69-305
3. P07: Thursday: 2-3pm: 47A-250

Homework Assignments


Homework as **R Markdown** document.

Week	Dates	Lecture	Tutorial	Assignment Due
1	25/2–01/3	✓	x	x
2	04/3–08/3	✓	✓	x
3	11/3–15/3	✓	x	✓
4	18/3–22/3	✓	✓	x
5	25/3–29/3	✓	x	✓
6	01/4–05/4	✓	✓	x
7	08/4–12/4	✓	✓	✓
8	15/4–18/4	✓	x	x
9	22/4–26/4	x	x	x
10	29/4–03/5	✓	✓	✓
11	07/5–10/4	✓	x	✓
12	13/5–17/5	✓	✓	x
13	20/5–24/5	x	x	✓
14	27/5–31/5	x	x	x

Lecture 2


1. Sample Space, Outcomes and Events
2. Probability
3. Conditional Probability and Independence
4. Birthday Problem (Monte Carlo Method)

Venn Diagram



Venn Diagram


Given sample space Ω and a set $B \subset A$. Highlight A^c in a Venn diagram.




"everything
but $A =$
 $= \Omega \setminus A$

Venn Diagram


Given sample space Ω and a set $B \subset A$ and C . Highlight $A \cap B \cap C$ in a Venn diagram.



$$\Rightarrow A \cap B \cap C = \emptyset$$




$$\Rightarrow A \cap B \cap C = C$$



$$\underline{A \cap B \cap C}$$

Venn Diagram

If $C \subset B$ and $B \subset A$, what can we say about the relation between C and A ?



i) $C \subset A$

ii) $A \cap B \cap C = C$


Operation Law - 1

$$\underline{(A \cup B) \cap C} = \underline{(A \cap C) \cup (B \cap C)}$$

LHS



RHS




Operation Law - 2

$$\underbrace{(A \cap B)}_{\text{orange}} \cup \underbrace{C}_{\text{orange}} = \underbrace{(A \cup C)}_{\text{orange}} \cap \underbrace{(B \cup C)}_{\text{pink}}$$

LHS



RHS




Operation Law - 3


(1st De Morgan's Law)

$$\overline{A \cup B} = \overline{A} \cap \overline{B}$$

LHS





RHS




Operation Law - 4

(2nd De Morgan's Law)

$$\overline{A \cap B} = \overline{A} \cup \overline{B}$$



Example - Network



Three unreliable systems

Left: Series system works only if all components work.

Center: Parallel system works if at least 1 component works.

Right: 2-out-of-3 system works if at least 2 components work.

Example - Network

Let A_i be the event that the i^{th} component works.

Express the event that “Series system” works using A_i .

$$\underline{A_1 \cap A_2 \cap A_3}$$

Express the event that “Parallel system” works using A_i .

$$\underline{A_1 \cup A_2 \cup A_3}$$

Express the event that “2-out-of-3 system” works using A_i .

$$\underline{(A_1 \cap A_2) \cup (A_1 \cap A_3) \cup (A_2 \cap A_3)}$$

Probability = Likelihood

Probability is a measure of how likely the outcome of an experiment is.

Probability = Likelihood

Probability is a measure of how likely the outcome of an experiment is.

Example:

Sample space Ω is the income of all aerospace engineers in Australia.

Let A be the event that the income of at least 90% of engineers is above 150K a year.

Probability = Likelihood

Probability is a measure of how likely the outcome of an experiment is.

Example:

Sample space Ω is the income of all aerospace engineers in Australia.

Let A be the event that the income of at least 90% of engineers is above 150K a year.

To measure the likelihood of this event A , we ask:

What is the probability/likelihood that at least 90% of engineers have an income of above 150K a year?

Probability - Formal definition

Definition

Let \mathcal{F} be the set of all subsets of Ω .

Probability - Formal definition

Definition

Let \mathcal{F} be the set of all subsets of Ω .

A probability $P : \mathcal{F} \rightarrow [0, 1]$ such that the following holds:

Probability - Formal definition

Definition

Let \mathcal{F} be the set of all subsets of Ω .

A probability $P : \mathcal{F} \rightarrow [0, 1]$ such that the following holds:

1. $0 \leq P(A) \leq 1$ for any $A \in \mathcal{F}$

Probability - Formal definition

Definition

Let \mathcal{F} be the set of all subsets of Ω .

A probability $P : \mathcal{F} \rightarrow [0, 1]$ such that the following holds:

1. $0 \leq P(A) \leq 1$ for any $A \in \mathcal{F}$
2. $P(\Omega) = 1$

Probability - Formal definition

Definition

Let \mathcal{F} be the set of all subsets of Ω .

A probability $P : \mathcal{F} \rightarrow [0, 1]$ such that the following holds:

1. $0 \leq P(A) \leq 1$ for any $A \in \mathcal{F}$
2. $P(\Omega) = 1$
3. If $A_1 \cap A_2 = \emptyset$ then $P(A_1 \cup A_2) = P(A_1) + P(A_2)$

Probability - Formal definition

Definition

Let \mathcal{F} be the set of all subsets of Ω .

A probability $P : \mathcal{F} \rightarrow [0, 1]$ such that the following holds:

1. $0 \leq P(A) \leq 1$ for any $A \in \mathcal{F}$
2. $P(\Omega) = 1$
3. If $A_1 \cap A_2 = \emptyset$ then $P(A_1 \cup A_2) = P(A_1) + P(A_2)$

Generalized:

If $A_i \cap A_j = \emptyset$ for all $i \neq j$, then

$$P(\cup_{i=1}^k A_i) = \sum_{i=1}^k P(A_i)$$

Probability - Properties

► $P(\emptyset) = \underline{0}$

► $P(\overline{A}) = \underline{1 - P(A)}$

► If $A \subset B$ then $P(A) \leq P(B)$

► $P(A \cup B) = \underline{P(A) + P(B) - P(A \cap B)}$

Probability - Discrete Sample Space

$$\Omega = \{a_1, a_2, \dots, a_n\},$$

Probability - Discrete Sample Space

$$\Omega = \{a_1, a_2, \dots, a_n\}, \text{ Event } A = \{a_1, a_4, a_6\}$$

Probability - Discrete Sample Space

$\Omega = \{a_1, a_2, \dots, a_n\}$, Event $A = \{a_1, a_4, a_6\}$, then

$$P(A) = P(a_1) + P(a_4) + P(a_6)$$

Probability - Discrete Sample Space

$\Omega = \{a_1, a_2, \dots, a_n\}$, Event $A = \{a_1, a_4, a_6\}$, then

$$P(A) = P(a_1) + P(a_4) + P(a_6)$$

In general:

$$P(A) = \sum_{i: a_i \in A} P(a_i)$$

Probability - Discrete Sample Space

$\Omega = \{a_1, a_2, \dots, a_n\}$, Event $A = \{a_1, a_4, a_6\}$, then

$$P(A) = P(a_1) + P(a_4) + P(a_6)$$

In general:

$$P(A) = \sum_{i: a_i \in A} P(a_i)$$

Example:

The probability that a machine produces at least 4 products an hour but less than 10 is

$$\underbrace{P(\text{machine produces 4 prod}) + P(\text{machine produces 5 prod}) + \dots + P(\text{machine produces 9 prod})}.$$

Probability

How do we get the values for $P(a_i)$?

- ▶ e.g. samples from experiments/available data collection
- ▶ **geometric probability**

Probability

How do we get the values for $P(a_i)$?

- ▶ e.g. samples from experiments/available data collection
- ▶ **geometric probability**

If all events are equally likely and Ω is finite, then

$$P(A) = \frac{|A|}{|\Omega|}$$

If Ω discrete, then $|A|$ = number of non-decomposable elements in set/event A .

If Ω continuous, then $|A|$ = length of the (continuous) interval representing A .

Example - Phone Number

If each digit from 0 to 9 of a 9-digit phone number are equally likely, what is the probability that:

- ▶ the phone number contains nine times the number 9?

$$P = \frac{|A|}{|\Omega|} = \frac{1}{10^9}$$

- ▶ the phone number contains exactly one 2?

$$P = \frac{|B|}{|\Omega|} = \frac{9 \cdot 9^8}{10^9}$$

for positions of the 2

- ▶ the phone number contains at least two 5?

$$P = 1 - P(\text{at most two 5}) = 1 - [P(\text{no 5}) + P(\text{one 5})]$$
$$= 1 - \left[\frac{9^9}{10^9} + \frac{9 \cdot 9^8}{10^9} \right]$$

for positions of the 5

Probability - Continuous Sample Space

geometric probability for continuous sample space

$\Omega = [a, b]$ or any other uncountable set (e.g. any subset of \mathbb{R})

Event $A = (a_1, b_1) \subset [a, b]$ and any value in Ω is **equally likely**

Probability - Continuous Sample Space

geometric probability for continuous sample space

$\Omega = [a, b]$ or any other uncountable set (e.g. any subset of \mathbb{R})

Event $A = (a_1, b_1) \subset [a, b]$ and any value in Ω is **equally likely** ,
then

$$P(A) = \frac{|A|}{|\Omega|} = \frac{b_1 - a_1}{b - a}$$


Example:

The probability that a randomly chosen number (not necessarily an integer) between 0 and 100 lies in $(0, 5)$ is

$$\underline{P(A) = \frac{5-0}{100-0} = \frac{1}{20} .}$$

Conditional Probability

Suppose we know that event A occurred. How does that affect the probability that B will occur?



\Rightarrow if A happened, then in the shaded area,
so B can only happen in $A \cap B$.

Conditional Probability

So, the relative change/likelihood that B occurs is

$$P(\text{ B happens, given that A happened}) = \frac{P(A \cap B)}{P(A)}$$

= “Probability that B occurs, given that A occurred”,
denoted by

$$P(B|A) = \frac{P(A \cap B)}{P(A)}$$

Conditional Probability

We therefore have

$$P(B|A) = \frac{P(A \cap B)}{P(A)}$$

multiply both sides by $P(A)$ to get (so-called "Chain Rule"):

Conditional Probability

We therefore have

$$P(B|A) = \frac{P(A \cap B)}{P(A)}$$

multiply both sides by $P(A)$ to get (so-called "Chain Rule"):

$$P(A \cap B) = P(B|A)P(A)$$

Conditional Probability - Example

Example:

Suppose you take two math courses: *Calculus for Engineers* and *Statistics*.

The event A , that the mean grade is 6, is then:

$$A = \{(\text{Calc} = 5, \text{Stat} = 7), (\text{Calc} = 6, \text{Stat} = 6), (\text{Calc} = 7, \text{Stat} = 5)\}.$$

Let B be the event that you had at least one 7.

$$\text{Then } A \cap B = \{(\text{Calc} = 5, \text{Stat} = 7), (\text{Calc} = 7, \text{Stat} = 5)\}.$$

Conditional Probability - Example

Example:

Suppose you take two math courses: *Calculus for Engineers* and *Statistics*.

The event A , that the mean grade is 6, is then:

$A = \{(\text{Calc} = 5, \text{Stat} = 7), (\text{Calc} = 6, \text{Stat} = 6), (\text{Calc} = 7, \text{Stat} = 5)\}$.

Let B be the event that you had at least one 7.

Then $A \cap B = \{(\text{Calc} = 5, \text{Stat} = 7), (\text{Calc} = 7, \text{Stat} = 5)\}$.

Given that the mean grade is 6, what is the probability that one grade was a 7?

Considering all cases are equally likely.

$$P(B|A) = \frac{P(A \cap B)}{P(A)} = \frac{|A \cap B|}{|A|} = \frac{2}{3} \quad |\Omega| = 7^2$$

Conditional Probability

Can we extend that result?

$$P(A \cap B \cap C) = \underline{P(C|A, B) \cdot P(B|A) \cdot P(A)}$$

Even more general:

$$P(\cap_{i=1}^k A_i) = P(A_k|A_1, \dots, A_{k-1}) \cdot P(A_{k-1}|A_{k-2}, \dots, A_1) \cdots \\ \cdots P(A_2|A_1)P(A_1)$$

Conditional Probability - Example

We draw 3 balls from a bowl with 6 white and 4 black balls.

Conditional Probability - Example

We draw 3 balls from a bowl with 6 white and 4 black balls.

What is the probability that all balls will be black if you do return them after every drawing?

$$\underline{P = \left(\frac{4}{10}\right)^3}$$

What is the probability that all balls will be black, if the draws are not returned?


$$\underline{P = \frac{4}{10} \cdot \frac{3}{9} \cdot \frac{2}{8}}$$

Law of Total Probability

Let B_1, B_2, \dots, B_n be a **partition** of Ω , that is
 $B_i \cap B_j = \emptyset$ for all $i \neq j$ and $\cup_{i=1}^n B_i = \Omega$

Then $P(A) = \sum_{i=1}^n P(A \cap B_i)$

and using conditional probabilities, we have



$$P(A) = P(A \cap B_1) + P(A \cap B_2) + \dots,$$

\neq \neq

$$P(A) = \sum_{i=1}^n P(A \cap B_i) = \sum_{i=1}^n P(A|B_i)P(B_i)$$

Law of Total Probability - Example

Semiconductor Failures:

P of Failure	Level of Contamination
0.10	High
0.05	Medium
0.01	Low

In a day, 15% of the chips are subjected to high levels of contamination, 25% to medium levels of contamination, and 60% to low levels of contamination.

F = “failure”, H = “High contamination”, M = “Medium contamination”, L = “Low contamination”. (So partition is the levels of contamination.)

$$\begin{aligned}P(F) &= P(F|H)P(H) + P(F|M)P(M) + P(F|L)P(L) \\&= 0.10 \cdot 0.15 + 0.05 \cdot 0.25 + 0.01 \cdot 0.60.\end{aligned}$$

Independence

Two events A and B are **independent** if any of the following holds:

Independence

Two events A and B are **independent** if any of the following holds:

► $P(A \cap B) = \underline{P(A)P(B)}$

Independence

Two events A and B are **independent** if any of the following holds:

- ▶ $P(A \cap B) = \underline{P(A)P(B)}$

- ▶ $P(A|B) = \underline{P(A)}$

Independence

Two events A and B are **independent** if any of the following holds:

▶ $P(A \cap B) = \underline{P(A)P(B)}$

▶ $P(A|B) = \underline{P(A)}$

▶ $P(B|A) = \underline{P(B)}$

Independence

Two events A and B are **independent** if any of the following holds:

▶ $P(A \cap B) = \underline{P(A)P(B)}$

▶ $P(A|B) = \underline{P(A)}$

▶ $P(B|A) = \underline{P(B)}$

This can be extended:

Events A_1, A_2, \dots, A_k are independent if

$$P(A_1 \cap A_2 \cap \dots \cap A_k) = P(A_1)P(A_2) \dots P(A_k) = \underline{\prod_{i=1}^k P(A_i)}$$